

Studies on Anthelmintic activity of *Ficus religiosa* fruits

Dr. Devendra N. Pandey and Dr. Sandeep Kumar Shukla

Department of Zoology
Govt. S.K.N.(P.G.) College, Mauganj Rewa M.P India
Govt. Maharaja (P.G.) College Chhatarpur M.P.India
Email- sandeepshukla910@gmail.com

Abstract:

In recent time use of plants natural products has increased tremendously, and there is a very demand of herbal products for therapeutic, clinical and agricultural purpose. In the present study aqueous extracts of fruits of commonly occurring plant of genus *Ficus religiosa* compared for their invitro anthelmintic activity in order to estimate the most potent of them. The extracts were found not only to paralyze (Vermifuge) but also to kill (Vermicidal). Results revealed that aqueous extract of fruits of *Ficus religiosa* is more potent in the test worms (*Pheretima Posthuma*).

Keywords: Anthelmintic activity *Ficus religiosa*, *Pheretima posthuma*.

Introduction:

Medicinal plants have served through the ages, as a constant source of medicaments for the exposure of variety of diseases. The history of herbal medicine is as old as human civilization. The plants are known to provide a rich source of botanical anthelmintics, antibacterials and insecticides Satyavati et. al (1976) and Lewis et. al. (1977).

There are more than 800 species and 2000 varieties of *Ficus* genus, most of which are native to old world tropics. *Ficus Religiosa* (Pipal tree) is some of the commonly occurring trees of this genus belonging to family *Moraceae* Mousa et. al. (1994). *F. religiosa* was reported to have anthelmintic potential. Stem

and bark extracts of *F. Religiosa* proved lethal to *Ascaridia galli* in vitro (Manoj et. al. 2008 and Kaushik et. al. 1981). Methanolic extract of bark of *F. Religiosa* was 100% lethal to *Haemonchus contortus* worms (Iqbal et. al.2001). The latex of some species of *Ficus* (*Moraceae*) i.e. *F. inspida* and *F. carica* was also reported to have anthelmintic activity De Amarin (1999). Based on this an attempt has been made to compare and evaluate the anthelmintic potency of aqueous extracts of fruits of *F. religiosa*.

Materials & Methods:

Plant Material

Fresh fruits of *F. religiosa* was collected from Mauganj Rewa district, Madhya Pradesh, India. These Plants material were authenticated at the Department of Zoology, Govt. S.K.N.(P.G.) College, Mauganj Rewa M.P.. These fresh fruits were then crushed and used to obtain aqueous extracts.

Preparation of Extract

200 G of fruits of *Ficus religiosa* were boiled separately with 1500 ml of double distilled water for 1h. Then they were kept at room temperature for 24h and then filtered through the muslin cloth. The filtrate then obtained was then concentrated to thick slurry and then residue was again boiled for 1h and filtered. The filtrate thus obtained was added to the thick slurry of first step. The resultant solutions thus obtained were boiled again to get thick concentrated extracts. These are then dried and used as powders. The

percentage yields were found to be 10.24% for *Ficus religiosa* respectively.

Experimental Animals

Indian adult earthworms (*Pheretima posthuma*) were used to study anthelmintic activity. The earthworms were collected from moist soil and washed with normal saline to remove all fecal matter. The earthworms of 3-5 cm in length and 0.1- 0.2 cm in width were used for all experimental protocol. The earthworm resembles both anatomically and physiologically to the intestinal roundworms parasites of the human being, hence can be used to study the anthelmintic activity (Nirmal et. al. 2007) .

Drugs and Chemicals

Albendazole (Glaxo Smithkline Pvt. Ltd.), Dimethyl Formamide, DMF, (Thomos Baker Chemicals Pvt. Ltd.) were used during the experimental protocol.


Anthelmintic Activity:

For the anthelmintic activity of aqueous extract of fruits of *F. religiosa*, Indian earthworms (*Pheretima Posthuma*) of 3-5 cm in length and 0.1-0.2 cm in width were used. The animals were divided in to five groups containing six earthworms in each group. All the extracts and standard drug solution were freshly prepared before starting the experiments. Extracts and the standard drug solution were freshly poured in different petridishes. All the earthworms were washed in normal saline before they were released into 10 ml of respective formulation as follows: Vehicle (5% DMF in normal Saline), Piperazine hydrate (37.5 mg/ml), aqueous extract (37.5 mg/ml) of *F. religiosa*. Observations were made for the time taken to paralyze (Paralysis was said to occur when the worms did not revive even in normal saline) and death (death was concluded when the worms lost their motility followed with their body colors

fading away). All the results were expressed as a mean + SEM of six animals in each group.

Result and Discussion:

Preliminary phytochemical analysis showed the presence of carbohydrates, flavanoids, aminoacids, steroids, saponins and tannins like phytoconstituents. Some of these phytoconstituents are responsible for anthelmintic activity. It is evident from the observations made in figure I *F. religiosa* had shown anthelmintic activity. Aqueous extract of fruits of *F. religiosa* shows paralysis at 21.20 minute and death at 39.42 minute. The standard drug Albendazole Shows paralysis at 2.35 minute and death at 5.20 minute.



Discussion:

It is evident from the findings of the present study that aqueous extracts of *F. religiosa*. was found to be least potent . Further studies using in vivo models are required to carry out and establish the effectiveness and pharmacological rationale. Moreover, phytochemical studies are also needed to lay down recommendation on scientific ground.

References:

1. Satyavati G.V., Raina M. K., Sharma M., Medicinal Plants of India., 1976, 1, 201–

206. Indian Council of Medical Research, New Delhi.
2. Lewis W.H., and Elvin-Lewis M.P.H., Medicinal Botany Plants Affecting Man's Health. John Wiley and Sons, 1977, New York.
3. The Wealth of India, Volume- (F-G), In: A dictionary of Indian Raw materials and industrial products. New Delhi, Council of Scientific and Industrial Research, 2005, 24-26.
4. Husain A., Virmani O P., Popli S P, Misra L N., Gupta M M., Srivastava G N., et al., Dictionary of Indian Medicinal Plants, Lucknow, India, CIMAP., 1992, 546.
5. Mousa O., Vuorela P., Kiviranta J., Wahab S A., Hiltohen R., Vuorela H., Bioactivity of certain Egyptian *Ficus species*, J Ethnopharmacol., 1994, 41, 71-76.
6. Manoj Aswar., Urmila Aswar., Bhagyashri Watkar., Minakshi Vyas., Akshaya Wagh., Kishore N Gujar., Anthelmintic activity of *Ficus benghaensis*, Int J Green Pharmacy., 2008, 170-172.
7. Kaushik R K., Katiyar J C., Sen A B., A new in vitro screening technique for anthelmintic activity using *Ascaridia galli* as a test parasite, Indian J Anim Sci., 1981, 51, 869-72.
8. Iqbal Zafar., Nadeem Qazi Khalid., Khan M N., Akhtar M S., Faisal Nouman Waraich, In vitro anthelmintic activity of *Allium sativum*, *Zingiber officinale*, *Cucurbita mexicana* and *Ficus religiosa*, Int J Agr Biol 2001., 3, 454-457.
9. De Amorin A., Borba H R., Carauta J P., Lopes D., Kaplan M A., Anthelmintic activity of the latex of *Ficus Species*, J Ethnopharmacol., 1999, 64, 255-258.
10. Nirmal S A., Malwadkar G., Laware R B., Anthelmintic activity of *Pongamia glabra*, Songklanakarin J Sci Technol., 2007, 29, 755-757.