

**DIVERSITY of MOSQUITO FAUNA in THREE  
SELECTED SITES of MADURAI DISTRICT,  
TAMILNADU, INDIA.**

K. RAMESH KUMAR and N. NATTUTHURAI

PG & RESEARCH DEPARTMENT OF ZOOLOGY

VIVEKANANDA COLLEGE

TIRUVEDAKAM WEST

MADURAI – 625234

TAMILNADU

Email: pnkramesh@gmail.com and

Pnkramesh29@yahoo.com

## ABSTRACT

**In the present study, the mosquito density of three selected sites of Madurai district viz., Sholavandan, Duraisamy nagar and Usilampatti was surveyed. The results of the study indicate the presence of 10 species of mosquitoes which are grouped under four genera namely *Aedes*, *Anopheles*, *Armigeres* and *Culex*. The numbers of mosquitoes collected belonging to various species were low with an exception of *Culex quinquefasciatus*. The diversity measures (Shannon's and Simpson's) shows slight variation among the three selected sites studied. The Shannon's index for Duraisamy nagar is slightly higher (2.008) when compared to Shoalvandan and Usilampatti (1.891 and 1.794). The Simpson's index for Duraisamy nagar is slightly higher (0.8258) when compared to Sholavandan and Usilampatti (0.7916 and 0.7568). In the study period, the maximum density of the mosquito population noted in the month of July and August. The minimum density of mosquito population obtained in the month of September.**

**Key words:** *Aedes*, *Anopheles*, *Armigeres* and *Culex*

## INTRODUCTION

Mosquito constitutes the most important single family of insects that affect the human health

everywhere. In spite of several attempts to control them, these remarkably adapted mosquitoes continue to successfully coexist with man, feeding on him and his domesticated animals (Pandian, 1998). Besides the blood loss, they are capable of transmitting many diseases like filariasis, malaria, yellow fever, Japanese encephalitis, dengue etc, (Dutta, *et al.*, 2003; Service, 1983). The voracious feeding habit high fecundity rate, dispersal potential and successful exploitation of environment are the causes for proliferation of these mosquitoes throughout the world (Pandian, 1990).

Mosquitoes are found in all types of environments associated with water such as sewage water, stagnant water, septic tanks etc. There is an urgent need to check the proliferation of population of vector and non-vector mosquitoes in order to reduce vector borne diseases and their nuisance by using appropriate control methods. The breeding habitat is crucial for mosquito population dynamics, because it is the location where many important life cycle processes take place. The processes are development of larva, emergence of adults, resting, swarming and mating of adults (Reuben, 1978).

As the density of vector and non vector has been increasing steadily, it is an essential to use all appropriate technological and management techniques to bring about an effective degree of

control in a cost effective manner (Kumar *et al.*, 2002). The management of mosquitoes requires to collect adequate knowledge about the species diversity and distribution pattern in a given area in order to evolve suitable strategy and to implement the same for the meaningful control of the population and in turn to reduce the menace and the incidence of the disease (Rajavel *et al.*, 2001).

Though several studies have been conducted at various places in the world, particularly in India and in a few places in Tamilnadu, there is a need to study the bionomics of mosquitoes in all areas. Hence, an attempt has been made to survey the mosquito fauna in three selected sites of Sholavandan, Duraisamy nagar and Usilampatti, Madurai district of Tamilnadu.

#### MATERIALS AND METHODS

##### *Sampling strategy:*

The mosquito sampling was carried out in three selected sites of Sholavandan, Duraisamy nagar and Usilampatti, Madurai district of Tamilnadu. Monthly sampling was done from July 2010 to December 2010. The Sampling was done between 6.00 pm to 6.00 am hours. The sampling sites were chosen at random and each sampling was done at the same place.

##### *Collection of adult female mosquitoes:*

The patterns of biting rhythm of mosquitoes were studied in two different sites in the selected

village by collecting the biting mosquitoes continuously throughout the night by using human bait. Therefore, human was considered as bait for the present study. The biting mosquitoes were collected by using transparent vials (4.5 x 2.5 cm) with a lid. The vials were directed towards the mosquitoes and placed suddenly over it, while they were biting. The collected mosquitoes were anaesthetized and killed by using ether and kept in separate vials, which were labeled with time and place of collection. The collected adult female mosquitoes were identified by the experts of Centre for Research in Medical Entomology (CRME) in Madurai.

##### *Diversity measures:*

The data collected in the present study was used in to calculate various diversity measures by following Southwood (1978).

#### RESULTS AND DISCUSSION

Mosquitoes had evolved by the Jurassic period, approximately 210 million years ago (Edwards, 1932). This was around the time continental drift began (Wilson, 1963), leading to the continental fragmentation and geographic isolation presumably promoting rapid speciation. There are more than 3200 species of mosquitoes belonging to 37 genera recorded so far world over and are grouped in to three subfamilies, Anophelinae, Culicinae and Toxorhynchitinae, which consist of 3 genera, 33 genera and 1 genus respectively (Ward, 1992). Out of

these, genus *Anopheles* consists of about 420 species, *Aedes* 950 species, *Culex* 800 species and *Mansonia* 25 species.

The voracious feeding habit, high reproductive and dispersal potential and successful exploitation of the environment are the causes of their rapid proliferation (Berlin, 1972). Moreover the distribution of mosquitoes is almost worldwide as they range from equator nearly to the poles and from sea level to atleast 7000 feet altitude.

Some species are important as vectors of malaria, filariasis, yellow fever, dengue fever and other arboviral diseases (Ross 1965, Service, 1983 and Rai, 1999). In India vector mosquitoes belong to four genera viz., *Anopheles* (malaria), *Aedes* (dengue fever), *Culex* (filariasis and Japanese encephalitis) and *Mansonia* (filariasis).

The Indian mosquito fauna include 255 species grouped under 16 genera. Fifty eight species belong to genus *Anopheles*, 57 species to *Culex*, 111 species to *Aedes* and 7 species to *Mansonia* (Nagpal and Sharma, 1995).

It has been reported that in Madurai there are 27 species of mosquitoes belonging to the genera *Aedes*, *Anopheles*, *Armigeres*, *Culex* and *Mansonia* (Pandian 1998). Among these genera of mosquitoes *Anopheles*, *Aedes* and *Culex* are considered to be most important because of their potential ability in

transmitting fatal diseases like Malaria, Dengue, Filariasis and Japanese encephalitis respectively.

In the present study, the survey conducted during the period of July to December shows that ten species belonging to four genera namely, *Aedes*, *Armigeres*, *Anopheles* and *Culex* are prevalent in the study area.

*Armigeres* is represented by only one species. *Anopheles* is represented by two species namely, *Anopheles culicifacies* and *Anopheles stephensi*. *Aedes* is represented by four species namely, *Ae. aegypti*, *Ae. albopictus*, *Ae. jamesi*, and *Ae. vexans*. *Culex* is represented by three species namely, *Cx. fuscocephala*, *Cx. bitaeniorhyncus* and *Cx. quinquefasciatus*. The numbers of mosquitoes collected belonging to various species were low with an exception of *Culex quinquefasciatus*.

Table 1. provides the various diversity measures of mosquito fauna recorded in Sholavandan during July 2010 to December 2010. The mosquito dominance was maximum in the month of September. The Shannon's index was maximum (1.891) in the month of July and minimum (1.307) in the month of September. A maximum of Simpson's index value (0.7916) was observed in July and a minimum index value (0.5441) was noted in the month of September.

The Dominance and diversity indices of mosquito fauna recorded in Duraisamy nagar are given in Table 2. The highest mosquito dominance was recorded in the month of September. The Shannon's index was maximum (2.008) in the month of October and minimum (1.461) in the month of September. The Simpson's index was maximum (0.8258) in the month of July and minimum (0.628) in the month of September.

Table 3 provides the various diversity measures of mosquito fauna recorded in Usilampatti during July 2010 to December 2010. The dominance of mosquito population was maximum in the month of September. High Shannon's and Simpson's diversity indices (1.794 and 0.7568) were calculated for the month of August minimum (1.324 and 0.5504) in the month of September.

It is well known that some of the mosquitoes which were originally zoophilic and sylvatic have adapted to feeding on humans and became peridomestic and even periurban due to deforestation. The involvement of man in certain host-parasite cycles will depend on the effect of his activities on the breeding sites of vectors, their capacity to adopt to new ecology and the presence of animal reservoirs as well as human behavior pattern.

In India, *Aedes albopictus* a vector of dengue haemorrhagic fever was a sylvatic species as found in the present survey. Of the study period, the month of August showed the high dense of all mosquito species followed by the November.

In the present study, the maximum density of the mosquito population noted in the month of July and October and the minimum density of mosquito population obtained in the month of September.

Biodiversity conservation needs the base line data of all fauna and flora, even the lesser known groups, as all have to play their own role as producers, consumers, pollinators and decomposers. The diverse micro and macro climatic conditions, harbors a variety of invertebrate fauna including mosquitoes that play a significant role in conserving and maintaining biodiversity in the study area.

#### ACKNOWLEDGEMENT

The authors greatly acknowledged the Secretary, Principal and Head of the Department of Zoology of Vivekananda College for providing facilities during the course of studies.

#### REFERENCES

- Dutta, P.S., Khan, S.A., Khan, A.M., Sharma, C.K., Hazirika, N.C. and Mahanta, J. (2003) Survey of medically important mosquito fauna in Mizoram. *Entomon* **28(3)**: 237-240.

- Kumar, S.R., Pandian, R.S. and ismail, M. (2002) Biting and breeding behaviour of mosquitoes in Devakottai, Tamilnadu, Southindia. *Env.Pol.Tech* **1** (2): pp 87-92.
- Pandian, R.S. (1990) The seasonal prevalence of adults of *Armigeres subalbatus* (Diptera: Culicidae) in Madurai, Tamilnadu. *J. Ecobiol* **2**: 172-174.
- Pandian, R.S. (1998) Biodiversity of mosquito fauna and efficacy of biopesticides against mosquitoes in an urban area in Tamilnadu. *Indian J. Environ. Sci.*, 2(1): pp 7-16.
- Rajavel, A.R., Munirathinam, A., Natarajan, R. and Vaidyanathan, K. (2001) species diversity of mosquitoes (Diptera: Culicidae) in mangrove ecosystem in South INDIA. *Entomon* **16** (3&4): 271-277.
- Reuban, R. (1978) A report on mosquitoes collected in the Krishna Godavari delta, Andhra Pradesh. *Indian J. Med. Res* **68**: 603-609.
- Southwood, T.R.E. (1978) Ecological methods. Chapman and Hall, U.K. pp 524.

Table: 1 Diversity indices of mosquito population in **Sholavandan**

<b>Diversity measures</b>	July 2010	Aug 2010	Sep 2010	Oct 2010	Nov 2010	Dec 2010
<b>Taxa S</b>	10	10	10	9	9	10
<b>Individuals</b>	224	277	523	443	555	481
<b>Dominance_D</b>	0.2084	0.2497	0.4559	0.2778	0.2819	0.275
<b>Shannon_H</b>	1.891	1.81	1.307	1.65	1.678	1.727
<b>Simpson_1-D</b>	0.7916	0.7503	0.5441	0.7222	0.7181	0.725
<b>Evenness_e<sup>H/S</sup></b>	0.6623	0.6108	0.3693	0.5783	0.595	0.5622

Table: 2 Diversity indices of mosquito population in **Duraisamy nagar**

<b>Diversity measures</b>	July 2010	Aug 2010	Sep 2010	Oct 2010	Nov 2010	Dec 2010
<b>Taxa_S</b>	10	10	9	10	10	10
<b>Individuals</b>	199	234	417	495	445	509
<b>Dominance_D</b>	0.1742	0.2161	0.372	0.2057	0.2577	0.2515
<b>Shannon_H</b>	2.008	1.886	1.461	1.936	1.782	1.791
<b>Simpson_1-D</b>	0.8258	0.7839	0.628	0.7943	0.7423	0.7485
<b>Evenness_e<sup>H/S</sup></b>	0.7445	0.6596	0.4791	0.6929	0.5941	0.5996

Table: 3 Diversity indices of mosquito population in **Usilampatti**

<b>Diversity measures</b>	July 2010	Aug 2010	Sep 2010	Oct 2010	Nov 2010	Dec 2010
<b>Taxa_S</b>	10	10	10	10	10	10
<b>Individuals</b>	248	332	580	500	561	578
<b>Dominance_D</b>	0.2623	0.2432	0.4496	0.2659	0.315	0.2704
<b>Shannon_H</b>	1.774	1.794	1.324	1.764	1.645	1.708
<b>Simpson_1-D</b>	0.7377	0.7568	0.5504	0.7341	0.685	0.7296
<b>Evenness_e<sup>H/S</sup></b>	0.5891	0.6014	0.3758	0.5837	0.5182	0.5517